On a problem of Hering concerning orthogonal covers of $\vec{K}_n$

A. Granville
University of Athens, GA

H.-D. O.F. Gronau
Universität Rostock

R.C. Mullin
University of Waterloo

April 18, 1997

Abstract

A Hering configuration of type k and order n is a factorization of the complete digraph $\vec{K}_n$ into n factors each of which consists of an isolated vertex and the edge-disjoint union of directed k-cycles, which has the additional property that for any pair of distinct factors say $\vec{G}_i$ and $\vec{G}_j$ there is precisely one pair of vertices, say $\{a, b\}$ such that $\vec{G}_i$ contains the directed edge (a, b) and $\vec{G}_j$ contains the directed edge (b, a). Clearly a necessary condition for a Hering configuration is $n \equiv 1 \pmod{k}$. It is shown here that for any fixed k, this condition is asymptotically sufficient, and, it is shown to be always sufficient for $k = 4$.

1 Introduction

Let $n \geq 1$ be an integer and let $\vec{K}_n$ denote the complete digraph on the n-element vertex set V. We consider collections $\vec{\mathcal{G}} = \{\vec{G}_1, \vec{G}_2, \ldots, \vec{G}_n\}$ of

spanning subdigraphs of $\vec{K}_n$. Note that the number of digraphs in $\vec{\mathcal{G}}$ coincides with the size of V. We call $\vec{\mathcal{G}}$ an **orthogonal cover** of $\vec{K}_n$ if

(i) every directed edge of $\vec{K}_n$ belongs to exactly one of the $\vec{G}_i$'s, and

(ii) for every two subdigraphs $\vec{G}_i$ and $\vec{G}_j (i \neq j)$ there is a unique pair $\{a, b\}$ of vertices such that $\vec{G}_i$ contains the directed edge (a, b) and $\vec{G}_j$ contains the directed edge (b, a).

Since the number of members of $\vec{\mathcal{G}}$ equals the number of vertices, we can use the vertex set V to index the members of $\vec{\mathcal{G}}$. Since each $\vec{G}_i, i \in V$ is spanning, we can consider the vertex in $\vec{G}_i$ to be distinguished, and we will refer to i as the root vertex (or simply the root) of $\vec{G}_i$. Then we refer to $\vec{G}_i$ as being the i**th page** of the cover. Furthermore, $\vec{G}_i$ is said to be **idempotent** if the vertex i occurs as an isolated vertex in $\vec{G}_i$. The cover $\vec{\mathcal{G}}$ is said to be idempotent if every page of $\vec{\mathcal{G}}$ is idempotent. Note that every page must have exactly $n - 1$ edges.

Hering [6] raised the question of determining, for a fixed integer $k \geq 3$, for which values of n does there exist an orthogonal cover of $\vec{K}_n$ in which every page consists of an isolated vertex and a vertex disjoint union of directed cycles of length k. Such a configuration will be called a Hering configuration of type k and order n.

Clearly a necessary condition for the existence of a Hering configuration of type k, defined on $\vec{K}_n$, is that $n \equiv 1 \pmod{k}$. Let $S_k = \{n$: there exists a Hering configuration of type k and order $n\}$. It will be shown that for each integer $k \geq 3$, there exists an integer N_k such that if $n \equiv 1 \pmod{k}$ and $n \geq N_k$, then $n \in S_k$.

2 Constructions

In this section we discuss two constructions for Hering configurations, one of which is direct, and the other recursive.

Theorem 2.1. *Let k be an integer, $k \geq 3$. Suppose that n is a prime power such that*
$n \equiv 1 (mod\ k)$. Then there exists a Hering configuration of type k and order n.

Proof. Let $GF(n)$ denote the finite field of order n. Since $n \equiv 1 \pmod{k}$, there exists an kth root of unity β in $GF(n)$. Define a quasigroup $Q = (GF(n), o)$ by $xoy = \beta x + (1-\beta)y$ for $x, y \in GF(n)$. Then Q is idempotent, that is $xox = x$, for all $x \in GF(n)$. Also

$$\underbrace{(\cdots((xoy)oy)\cdots)oy)}_{k \text{ times}} = \underbrace{\beta(\beta\cdots\beta x + (1-\beta)y) + (1-\beta)y)\cdots) + (1-\beta)y}_{k \text{ times}}$$

$$= \beta^k x + (\beta^{k-1} + \beta^{k-2} + \cdots + \beta + 1)(1-\beta)y = \beta^k x + (1-\beta^k)y = x,$$

since β is a k'th root of unity. Further, for a given pair of elements x and y, there exists a unique element u such that $(uox)oy = u$. Given these facts, it is easily verified that by defining $\vec{G}_i$ by

$$\vec{G}_i = \{(i)\} \cup \bigcup_{\substack{x \in GF(n) \\ x \neq i}} \{(x,\ xoi)\}$$

for $i \in GF(n)$, a Hering configuration of type k and order n is obtained. □

For the recursive construction, the notion of pairwise balanced design (PBD) is required. Let v be a positive integer, and K a subset of the positive integers. Then a pairwise balanced design $PBD[v, K]$ is a pair $(V, \mathcal{F})$ where V is a v-set and $\mathcal{F}$ is a family of subsets (called blocks) of v which satisfies the following:

(i) every pair of distinct elements of V occur in precisely one block;

(ii) the cardinality (size) of every block lies in K.

A set S of positive integers is said to be PBD-closed if it has the property that the existence of a PBD$[v, S]$ implies that v lies in S. A well known theorem of R.M. Wilson [9] states that a PBD-closed set S is ultimately periodic with period $\alpha = GCD\{s(s-1) : s \in S\}$.

Theorem 2.2. *Let k be a fixed integer ≥ 3. Then the set $S_k = \{n$: there exists a Hering configuration of type k and order $n\}$ is PBD-closed.*

Proof. It is shown in [4] that if there exists a PBD $[n, S]$ and for each $s \in S$ there exists an idempotent orthogonal covering of $\vec{K}_s$, then there exists an

idempotent orthogonal covering of $\vec{K}_n$ whose pages each consist of the idempotent together with vertex disjoint unions of the connected components, apart from the idempotents, of the pages of the covering $\vec{K}_s$. In the case at hand, these components are all directed cyles of length k, so the resulting configuration is a Hering configuration of type k and order n. Therefore S_k is PBD-closed. $\Box$

3 Asymptotic results on the sets S_k

In this section, we show that for fixed $k \geq 3$, there exists an integer N_k such that if
$n \equiv 1 \pmod{k}$ and $n \geq N_k$, then there exists a Hering configuration of type k and order n. To this end, let k be any integer, $k \geq 2$, and let $P(k) = \{p : p \text{ is a prime, } p \equiv 1 \pmod{k}\}$, and let $Q(k) = \{q : q = p^t, p \text{ is a prime, } t \text{ is a positive integer, } q \equiv 1 \pmod{k}\}$. For any non-empty set of positive integers S let $\beta(S) = GCD\{s(s-1) : s \in S\}$.

Lemma 3.1 *Let k be an integer, $k \geq 2$. Then there exist two primes p_1 and p_2 in $P(k)$ such that $GCD(p_1(p_1-1), p_2(p_2-1)) = 2k$.*

Proof. By Dirichlet's Theorem on primes in an arithmetic progression (see [1]), there exists a prime p_1 such that $p_1 \equiv 1 \pmod{2k}$, say $p_1 = 1 + 2ka$ for some positive integer a. Further by Dirichlet's theorem there exists a prime p_2 such that $p_2 \equiv 1 + 2k \pmod{2kp_1a}$, say $p_2 = 1 + 2k + 2kp_1ab$ for some positive integer b. Note that $p_2 > p_1$. Therefore

$$\begin{aligned} GCD(p_1(p_1-1),\ p_2(p_2-1)) &= GCD(p_1(p_1-1),\ p_2-1) \\ &= GCD(p_1 2ka,\ 2k + 2kp_1ab) \\ &= 2k\ GCD(p_1a,\ 1 + p_1ab) \\ &= 2k, \end{aligned}$$

as required. $\Box$

Corollary 3.1.1. *Let k be an integer, $k \geq 3$. Then*

(i) *$\beta(P(k))$ divides $2k$.*

(ii) *Further if k is even, then $\beta(P(k)) = k$.*

Proof. Part (i) is a direct consequence of Theorem 3.1 and the definition of $\beta(P(k))$. For part (ii), assume that k is even, that is, $k = 2s$ where $s > 2$. By Theorem 3.1, there exist primes p_1 and p_2 in $P(s)$ such that $GCD(p_1(p_1-1),\ p_2(p_2-1)) = 2s = k$. But p_1 and p_2 are both odd and they are both relatively prime to s, since they lie in $P(s)$. Hence $2s$ divides both $p_1 - 1$ and $p_2 - 1$, so p_1 and p_2 are in $P(k)$.
These are then the required primes. □

These results can be applied to the sets S_k as follows.

Theorem 3.2. *Let k be any integer, $k \geq 3$. Then there exists a constant N_k such that if $n \geq N_k$ and $n \equiv 1$ (mod k), then $n \in S_k$.*

Proof. By Theorem 2.1, we have $P(k) \subset Q(k) \subset S_k$, so S_k is non-empty and $k \leq \beta(S_k) \leq 2k$, with $\beta(S_k) = k$ if k is even.

Wilson's theory states that the PBD-closed set S_k is ultimately periodic with period $\beta(S_k)$, and that for any m in S_k there exists a constant C_m such that if $n \geq C_m$ and if $n \equiv m \pmod{\beta(S_k)}$, then $n \in S_k$. We consider two cases, namely k odd and k even. Suppose first that k is even. Then $\beta(S_k) = k$, and since S_k contains some member $m \equiv 1 \bmod k$, then by Wilson's theorem there exists a constant N_k as in the enunciation of this theorem.

Now consider the case when k is odd. Then the argument is slightly more difficult, since in this case $\beta(S_k) = 2k$. Now suppose that $m \equiv 1 \bmod k$. If m is odd, then $m \equiv 1 \pmod{2k}$, and if m is even, then $m \equiv k+1 \pmod{2k}$. Therefore if we can show that S_k contains both odd and even integers, an argument similar to that above applied to each of these cases will establish the existence of the required integer N_k. But since k is odd, then $2^{\phi(k)} \equiv 1 \pmod{k}$ where $\phi(k)$ is the Euler phi function, so $2^{\phi(k)} \in Q(k)$. Therefore S_k contains both odd and even integers, and the theorem follows. □

4 The spectrum of Hering configurations of type 4

Ganter and Gronau [2] have shown that $S_3 = \{n : n \geq 4, n \equiv 1 \pmod 3, n \neq 10\}$. To obtain an analogous result for S_4, we require the notion of the closure of a set of positive integers. Let K be any nonempty set of positive integers.

Then $B[K] = \{n : \text{there exists a } PBD[n, K]\}$ is clearly PBD closed and is called the closure of K.

It is shown in [5] and [7] that $B[\{5, 9, 13, 17, 29, 33\}] = \{n : n \geq 5, n \equiv 1(\text{mod } 4)\}$. But $\{5, 9, 13, 17, 29\}$ is a set of prime powers, and a Hering configuration of type 4 and order 33 is exhibited in table 1. Therefore $S_4 = \{n : n \geq 5, n \equiv 1(\text{mod } 4)\}$.

A Hering configuration of type 4 and order 33

$\{(0) \quad (1,2,4,3)\ (5,8,12,19)\ (6,28,20,16)\ (7,31,15,25)$
$(9,22,10,30)\ (11,27,24,17)\ (13,21,26,32)\ (14,23,18,29)\}\ (\text{mod } 33)$

TABLE 1

For $k = 5$, an exhaustive search shows that there is no Hering configuration of type 5 and order 6. However such configurations exist for $n = 11$ and 16 by theorem 2.1. Examples of Hering configurations of type 5 and orders 21 and 26 are exhibited in Table 2. However, with present methods a complete determination of $H(5)$ appears to be well beyond reach.

A Hering configuration of type 5 and order 21

$\{(0) \quad (1,2,4,7,12)\ (3,18,13,19,16)\ (5,17,10,14,6)$
$(8,15,11,9,20)\}\ (\text{mod } 21)$

A Hering configuration of type 5 and order 26

$\{((0,0)),\ ((1,0),\ (2,0)\ (4,0)\ (3,0),\ (0,1)),\ ((5,0),\ (8,0)\ (6,0),\ (10,0),\ (1,1)),$
$((7,0),\ (2,1),\ (11,0)\ (12,1),\ (7,1)),\ ((9,0),\ (8,1)\ (4,1),\ (9,1)\ (11,1)),$
$((12,0),\ (6,1)\ (5,1),\ (3,1)\ (10,1))\}$

$\{((0,1)),\ ((0,0)\ (5,0)\ (12,0)\ (7,0)\ (7,1)),\ ((11,0),\ (4,0),\ (1,0),\ (10,0),\ (3,1)),$
$((6,0),\ (4,1),\ (3,0),\ (8,1),\ (9,1)),\ ((8,0),\ (10,1)\ (2,0),\ (11,1),\ (1,1)),$
$((9,0),\ (12,1),\ (5,1),\ (2,1),\ (6,1))\}\ (\text{mod } 13, -).$

TABLE 2

The case of Hering configurations of type 6 is much more complete because of the fact that so many early members of S_6 are primes and prime powers. Let $N(6) = \{n : n \geq 7,\ n \equiv 1 (\mathrm{mod}\ 6)\}$, and $C(6) = B[Q(6)]$. It is shown in [8] and [10] that $C(6) \supseteq N(6) \backslash E$ where $E = \{55, 115, 145, 205, 235, 253, 265, 295, 319, 355, 391, 41$ $697, 745, 781, 799, 805, 1243, 1255, 1315, 1585, 1819, 1921\}$.

This result was improved by Greig [3] who showed that $\{295, 655, 1243, 1255, 1795, 1819,$ $1921\} \subset C(6)$. Therefore there exists a Hering configuration of type 6 for all $n \in N(6)$ with the possible exception of $n \in \{55, 115, 145, 205, 235, 253, 265, 319, 355, 391, 415, 44$ $649, 667, 685, 697, 745, 781, 799, 805, 1315, 1585, 1795\}$.

References

[1] L.E. Dickson, Theory of Numbers, Chelsea Press Vol. 1 (1971), 415.

[2] B. Ganter and H.-D. O.F. Gronau, On two conjectures of Demetrovics, Füredi and Katona on partitions, Discrete Math 88(1991), 149-155.

[3] M. Greig, Designs from projective planes and PBD bases, and designs from configurations in projective planes. Preprint.

[4] H.-D. O.F. Gronau, R.C. Mullin and P.J. Schellenberg, On orthogonal double covers of K_n and a conjecture of Chung and West, Journal of Combinatorial Designs (accepted).

[5] A.M. Hamel, W.H. Mills, R.C. Mullin, Rolf Rees, D.R. Stinson and Jainxing Yin, The spectrum of $PBD(\{5, k^*\}, v)$ for $k = 9, 13$, Ars Combinatoria 36(1993), 7-26.

[6] F. Hering, Balanced pairs, Annals of Discrete Math 20(1984), 177-182.

[7] E.R. Lamken, W.H. Mills and R.M. Wilson, Four pairwise balanced designs, Designs, Codes and Cryptography 1(1991), 63-68.

[8] R.C. Mullin and D.R. Stinson, Pairwise balanced designs with block sizes $6t + 1$, Graphs and Combinatorics 3(1987), 365-377.

[9] R.M. Wilson, An existence theory for designs II. The structure of PBD-closed sets and the existence conjectures, J. Combinatorial Theory 13(1972) 246-273.

[10] Zhu Lie and Chen Demeng, Orthogonal Steiner triple systems of order $6m+1$. (Unpublished manuscript).